

# Simultaneous magma and gas eruptions at three volcanoes in southern Italy: An earthquake trigger?

T.R. Walter<sup>1\*</sup>, R. Wang<sup>1</sup>, V. Acocella<sup>2</sup>, M. Neri<sup>3</sup>, H. Grosse<sup>1</sup>, J. Zschau<sup>1</sup>

<sup>1</sup>Department 2: Physics of the Earth, Helmholtz Centre Potsdam, GFZ German Research Centre for Geoscience, Telegrafenberg, 14473 Potsdam, Germany

<sup>2</sup>Dipartimento Scienze Geologiche, Università Roma Tre, Largo S.L. Murialdo 1, 00146 Rome, Italy

<sup>3</sup>Istituto Nazionale di Geofisica e Vulcanologica (INGV), Catania, Piazza Roma 2, 95123 Catania, Italy


## ABSTRACT

In September 2002, a series of tectonic earthquakes occurred north of Sicily, Italy, followed by three events of volcanic unrest within 150 km. On 28 October 2002, Mount Etna erupted; on 3 November 2002, submarine degassing occurred near Panarea Island; and on 28 December 2002, Stromboli Island erupted. All of these events were considered unusual: the Mount Etna northeast-rift eruption was the largest in 55 yr; the Panarea degassing was one of the strongest ever detected there; and the Stromboli eruption, which produced a landslide and tsunami, was the largest effusive eruption in 17 yr. Here we investigate the synchronous occurrence of these clustered events, and develop a possible explanatory model. We compute short-term earthquake-induced dynamic strain changes and compare them to long-term tectonic effects. Results suggest that the earthquake-induced strain changes exceeded annual tectonic strains by at least an order of magnitude. This agitation occurred in seconds, and may have induced fluid and gas pressure migration within the already active hydrothermal and magmatic systems.

## INTRODUCTION

Volcanoes interact with their environment on different scales, and with different modes and processes, ranging from climate and tidal relationships to tectonic interactions. Tectonic interactions, in particular, have received special attention in recent years. A correlation of earthquakes and eruptions was revealed by a statistical examination of global data catalogues, and these relationships may occur over distances exceeding hundreds of kilometers (Linde and Sacks, 1998). A mechanical relationship between apparently interlinked processes is, however, still not understood, partly due to the limited number of studied cases. Recent papers suggest that dynamic strain, together with long-term tectonic extension (Hill, 2008) and/or short-term static extension associated with earthquakes (Walter and Amelung, 2007), may increase the number of volcanic eruptions. A series of volcanic unrests occurred in 2002 in southern Italy in the weeks following an earthquake, and may help us to better understand such clustered events.

The Aeolian Arc is associated with the northwestward-subducting Ionian slab and currently hosts several active volcanoes. Tectonic deformation within the arc is heterogeneous, being subject to extensional tectonics in the east (including the volcanoes of Stromboli and Panarea), dextral shear tectonics in the center (including Vulcano and Lipari), and compressional tectonics in the west (Alicudi and Filicudi, Fig. 1; De Astis et al., 2003). The dextral shear



**Figure 1. Shaded relief map of Sicily and Aeolian Islands. Tectonic strain orientation: black arrows show compression in western volcanic Aeolian Arc, white arrows show extension in eastern Aeolian Arc (after Billi et al., 2007). Circles are earthquake magnitude  $M > 1$  epicenters from Advanced National Seismic System earthquake catalogue 2000–2005 (<http://earthquake.usgs.gov/anss/>), stars are mainshocks  $M > 5$ , Harvard centroid moment tensor solution is provided for Palermo  $M = 5.9$  event. Note that four  $M > 5$  earthquakes occurred, but two had depths  $>200$  km (17 May 2001 [ $M = 5.2$ ] and 5 May 2004 [ $M = 5.5$ ]), while September 2002 earthquake and its aftershocks were shallow ( $<30$  km). Earthquake main shocks and remotely triggered volcanoes investigated herein are shown with red symbols. Tectonic lines are from Billi et al. (2007). Ust—Ustica; Ali—Alicudi; Fil—Filicudi; S—Salina; Li—Lipari; V—Vulcano; Pan—Panarea; Str—Stromboli. Seismic stations at TOR (Tortorici) and AIO (Antillo).**

of the central Aeolian Arc is associated with a N-NW-S-SE-trending structure probably constituting the northernmost part of the Maltese Escarpment. This is the surface expression of a tear separating the subducting oceanic lithosphere (to the east) from the colliding continental lithosphere (to the west), and may facilitate the

extension and the rise of asthenospheric material at Mount Etna (Gvirtzman and Nur, 1999).

While there is a long-term interdependence between the southern Italy volcanism and tectonics (Neri et al., 1996; Lanzafame and Bousquet, 1997), the short-term link is still debated. Historical records suggest that some particular events

\*E-mail: twalter@gfz-potsdam.de.

of volcanism at Mount Etna have occurred in temporal proximity to large tectonic earthquakes (Feuillet et al., 2006). Synchronous activity at several volcanoes and their possible link to large tectonic earthquakes, however, has not been elaborated. In 2002, a significant earthquake took place west of the Aeolian Islands and was followed by a widespread aftershock sequence, and by major eruptions at Mount Etna and Stromboli Island and anomalous degassing at Panarea. Through observation, seismic investigation, and numerical modeling, this group of events is investigated here. Our study suggests that the volcanoes were further activated by dynamic pressure fluctuations associated with the earthquake, having implications that are important for understanding clustered activity and hazards in southern Italy and elsewhere.


## CHRONOLOGY OF THE EVENTS

### Palermo Earthquake

On 6 September 2002, at 01:21:27 (UTC), an earthquake occurred ~40 km northeast of Palermo, Sicily (Rovelli et al., 2004), and 130–140 km west of the very active volcanoes, Mount Etna and Stromboli Island (Fig. 1). The earthquake was followed by more than 600 aftershocks of  $M > 1$ , with hypocenters aligned in a northeastern continuation along an ~100 km segment of a  $050^\circ$  trending fault. The main shock killed two people, damaged several buildings in the Palermo area, and was felt in eastern Sicily in the cities of Catania and Messina. In northern Sicily, the earthquake is thought to have triggered the Cerda landslide (Agnesi et al., 2005) and affected physical parameter recordings at thermal springs (Caracausi et al., 2002). Seismological characteristics were detailed in the International Seismic Catalogue (ISC; <http://www.isc.ac.uk>), with the hypocenter located at  $38.36^\circ\text{N}$ ,  $13.69^\circ\text{E}$  at 12 km depth. The U.S. Geological Survey National Earthquake Information Center (NEIC) and Harvard seismology (HRVD) solutions provided a magnitude  $M_w = 5.9$ , with a focal mechanism nodal plane striking SW-NE (NEIC 242/60/145 or 351/60/35, and HRVD 26/50/40 or 267/60/133). As illustrated in Figure 2, the earthquake was soon followed by eruptions at Mount Etna and Stromboli Island, and degassing close to the island of Panarea. At these three volcanic centers, the change and scale of activity were very unusual, as detailed below.

### Mount Etna Eruption

Mount Etna erupted in July–August 2001 (Allard et al., 2006, and references therein). From 27 October 2002 to 28 January 2003, the volcano erupted again, associated with the opening of the northeast rift. This was the first northeast rift fissure eruption after 55 yr of quiescence (since 1947). Associated with this



**Figure 2. Volcanic activity (red lines) shortly after largest earthquake and its aftershocks in September 2002. Same earthquake data as in Figure 1; magnitudes >5 are indicated by stars. a—Mount Etna; b—Panarea; c—Stromboli.**

event, a major part of the east flank of the volcano was displaced eastward by meters, with surface fracturing and hazardous earthquakes; these events had not been observed in the previous decades (Neri et al., 2005).

### Panarea Degassing

Panarea is in the eastern Aeolian Arc and had its main period of activity in the Holocene. A constructional phase occurred between 150 and 105 ka, followed by discrete explosive eruptions until 8 ka, associated with slight submergence (Lucchi et al., 2006). Only reports of degassing activity are found for historical time. No dramatic increase or decrease in gas flux was ever instrumentally recorded before 3 November 2002 (Esposito et al., 2006). The anomalous period of increased degassing activity ended in January 2003. Observations showed that gas discharge occurred in at least three distinct areas ~3 km offshore of Panarea Island (BGVN 27:10, at <http://www.volcano.si.edu>). Geochemical monitoring revealed a dynamic behavior changing in time, space, and flux, with a component of the gases being directly associated with magmatic fluids (Capaccioni et al., 2007). The only other account of a similarly strong degassing episode refers to the year 1865 (Billi and Funicello, 2008).

### Stromboli Island Eruption

Stromboli Island, in the eastern Aeolian Arc, hosts one of the most active volcanoes with continuous archetypal Strombolian eruptions. These are usually associated with gas bubble rise, coalescence, and slug bursts, rather than juvenile effusion. Continuous radon gas measurements showed that summit degassing increased shortly after the 2002 Palermo earthquake (Cigolini et al., 2007). On 28 December 2002, Stromboli Island had its first dike-fed effusive eruption in 17 yr (since 1985), culmi-

nating two days thereafter in failure of part of the northern flank into the sea and the formation of a tsunami. The familiar Strombolian activity resumed in mid-2003 (Ripepe et al., 2005).

## STRESS AND STRAIN TRANSFER MODELING


Our calculations take the static and dynamic transfer of strain due to the 6 September 2002 Palermo earthquake into account. Strain changes were first calculated by producing synthetic seismograms at sites where we had actual seismic data available. We used seismic data from the Mediterranean Very Broadband Seismographic Network (MedNet <http://mednet.rm.ingv.it>) at a station in Antillo (AIO,  $37.9712^\circ\text{N}$ ,  $15.2330^\circ\text{E}$ , height = 751 m), recorded by a STS2 station at 20 Hz, and from another station maintained by INGV (Istituto Nazionale di Geofisica e Vulcanologica) Catania in Tortorici (TORT,  $38.040^\circ\text{N}$ ,  $14.810^\circ\text{E}$ , height = 540 m), recorded at 200 Hz by an accelerometer station (details are provided in the GSA Data Repository<sup>1</sup>). Stations AIO and TORT are located 142 and 104 km from the Palermo earthquake epicenter, respectively, and span the distance range of the investigated volcanoes (Mount Etna, 134 km; Panarea, 124 km; Stromboli Island, 141 km). Upon successful reproduction of the amplitudes of these seismograms in computer models, we were able to simulate strain changes at any other location, namely at Mount Etna ( $37.734^\circ\text{N}$ ,  $15.004^\circ\text{E}$ ), at Panarea ( $38.63^\circ\text{N}$ ,  $15.07^\circ\text{E}$ ), and at Stromboli Island ( $38.789^\circ\text{N}$ ,  $15.213^\circ\text{E}$ ). Because we were interested in how the earthquake caused transient changes at magma reservoirs, we first simulated

<sup>1</sup>GSA Data Repository item 2009065, details of model parameters, earthquake distribution, and waveform recordings, is available online at [www.geosociety.org/pubs/ft2009.htm](http://www.geosociety.org/pubs/ft2009.htm), or on request from [editing@geosociety.org](mailto:editing@geosociety.org) or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.

the seismic wave fields and then calculated the associated pressure fluctuation at depth (see the Data Repository). This provided a quantitative estimate of the scale of transient pressure changes under each of the volcanic centers, i.e., 2 km below sea level at Mount Etna, Panarea, and Stromboli Island. We assume that this is a reasonable depth that may host hydrothermal as well as shallow magmatic reservoirs.

The Palermo earthquake synthetic wave propagation essentially depends upon the earthquake source and strength considered (for model fault data, see Item DR1 in the Data Repository). This plane is discretized by 100 point sources, which are triggered by the rupture front propagating circularly from the hypocenter at a constant velocity of 2 km/s. The seismic moment of each point source is released via a set of Brune's (1970) subevents. We did not attempt to perfectly match the waveforms, but rather the three component amplitudes that yield information about the magnitude of induced dynamic strain changes. The characteristic duration of each point source is comparable to the rise time of the earthquake, which is related empirically to the magnitude and stress drop (Boore, 1983). Using the semi-analytical code by Wang (1999) to calculate synthetic seismograms, we produced the Green's functions for the standard seismic reference model IASPEI91. Synthetic seismograms of the earthquake are obtained by convolution between the Green's functions and the source functions described above.

The calculations show a large fluctuation of the three components at 2 km depth. As shown in Figure 3, the east and north components, as well as the vertical components of all synthetic waveforms, display 15 to ~18 s of time lag between P- and S-wave arrivals, which is consistent with the distance of 125–140 km between the earthquake hypocenter and the volcano locations. Amplitudes at all sites are similar to the true recordings at Antillo and Tortorici. At the Mount Etna site the vertical component is larger, while at the Panarea and Stromboli sites the north-south component is larger, which is related to the moment tensor solution applied in the initial rupture model and considered to be realistic. From these three components, we infer that the pressure changes are fluctuating for ~25–30 s at  $\pm 10$  kPa at Mount Etna, and  $\pm 8$  kPa at Panarea and Stromboli Island. Thus, the dynamic pressure fluctuations reach ~20 kPa and then fall back to near zero after the seismic waves pass. A small offset from the zero line is found due to static offset related to the permanent dislocation induced by the earthquake model. The static offsets are negligible (<1 kPa) and are thus an implausible eruption trigger, while the dynamic fluctuations (~20 kPa) exceed values known to have induced seismicity or volcanic activity elsewhere, as discussed below.



**Figure 3. Synthetic seismograms showing east-west (E), north-south (N), and vertical components (Z). From these three components, we determined the pressure change (P, shown in red) as function of time. Pressure fluctuations reached  $\pm 10$  kPa for Mount Etna and  $\pm 8$  kPa for the Panarea and Stromboli Island volcanoes.**


## DISCUSSION AND CONCLUSIONS

The observed degassing activity at Panarea and the types and scales of the eruptions at Mount Etna and Stromboli Island were each considered peculiar within the context of their recent decades of activity. Even more remarkable is the synchronous occasion of these events. Large eruptions and degassing events have been observed at the three volcanic centers in the past (e.g., in 1865), although never instrumentally recorded in such a close temporal proximity as in the period from October–December 2002. We investigated here the influence of a possible external trigger. As shown, the Palermo earthquake induced transient changes at the magmatic and hydrothermal systems of these volcanoes.

Alternatively, one may speculate whether the 2002 synchronous activity was an expression of a general geodynamic reorganization affecting the southern Tyrrhenian area. A geodynamic reorganization can cause static stress changes and thus act as a regional tectonic trigger, and may have locally led to both the Palermo earthquake and the simultaneous volcanic activity. However, regional seismicity does not suggest major plate movement (Item DR2). As suggested by Cigolini et al. (2007), and as quantitatively tested in this work, the possibility that the volcanic activity increased due to dynamic stress changes directly associated with the earthquake mainshock alone appears to be reasonable. Although the models presented herein are simplified and ignore complex heterogeneities and time-dependent rheology, they may help us to understand the simultaneous 2002 volcanic activity in Italy.

Model calculations suggest that pressure fluctuations of ~20 kPa occurred surrounding the magmatic and hydrothermal reservoirs of the volcanoes. These values may appear small

considering absolute pressures at magma stagnation levels (GPa), or magmatic overpressures required for magma chamber wall rupture and dike propagation (MPa). However, values of tens of kilopascals appear large in comparison to long-term plate tectonic forcing and to short-term extrinsic forcings, including various types of tidal and earthquake triggers. Long-term tectonic strain rates in the Aeolian Arc and at Mount Etna are generally <100 nanostrain  $\text{yr}^{-1}$  (D'Agostino and Selvaggi, 2004), which is about one order of magnitude smaller than values estimated from the Palermo earthquake model presented in this paper (Fig. 4). Dynamic triggering elsewhere even suggests that stress changes below those calculated in this work might be significant. For example, seismicity increases at the Long Valley caldera, California, associated with regional and teleseismic tectonic earthquakes were found to be triggered



**Figure 4. Conceptual model of time ( $t$ )-strain ( $\epsilon$ ) changes in the Panarea-Stromboli-Etna systems. Extensional tectonic strain built up in the long term, locally causing elevated pore pressure at hydrothermal and magmatic centers. Earthquake-induced short-term fluctuations exceeded the long-term signals. In the 2002 events, strain changes are similar to 10–20 yr of tectonic strain, but occurred within seconds.**

if the 5 kPa threshold was reached (Brodsky and Prejean, 2005). Such small changes may lead to a chain of adjustments within a magma-hydrothermal system already in a critical state and may explain the delay between the earthquake and observed volcanic activity. The chain of adjustments may begin with the excitation and ascent of gas bubbles, and associated pressure and density changes within a magmatic reservoir and other fluid-filled structures (Manga and Brodsky, 2006). Similar dynamic interaction may have occurred before, as in 1865, when strong degassing was observed at Panarea and eruptions occurred at Stromboli and Mount Etna following strong earthquakes.

We note that the volcanoes located within the eastern Aeolian Arc became active, while other volcanoes on the central and western arc did not show any significant changes. The structural tectonic configuration reveals that the eastern Aeolian Arc is subject to extensional tectonic strain, being strike-slip or transtensional in the central arc and compressional in the western arc (De Astis et al., 2003). Regions near Mount Etna are in part also subject to extensional tectonic strain (D'Agostino and Selvaggi, 2004) and to a complex local volcano-tectonic deformation additionally related to intrusions and gravitational spreading (Feuillet et al., 2006). In a simplistic view, this work may imply that volcanoes located in extensional tectonic environments are more prone to being activated by dynamic effects, providing a possible explanation of why the volcanoes closer to the Palermo earthquake did not show any response related to remote triggering. Although in the present scenario volcanoes located closer to the earthquake source are generally less active, such an interpretation is consistent with recent findings regarding the triggering of earthquakes (Hill, 2008) that suggest that extensional tectonic regimes are more vulnerable to dynamic triggering than compressional regimes. The fact that the synchronously excited volcanoes have already been in a near critical state, in terms of their magmatic systems (eruptions at Stromboli Island and Mount Etna) and hydrothermal system (Panarea), may be of additional importance, as the local pore pressures may have been already elevated when the remote trigger occurred.

#### ACKNOWLEDGMENTS

Three-component seismic data were generously provided by Antonio Rovelli; thanks to Matteo Picozzi and Domenico Di Giacomo for help with the data analysis. We also thank three *Geology* reviewers for their very constructive comments. This study was partly funded by the Deutsche Forschungsgemeinschaft (WA 1642/1-4), and Protezione Civile, project INGV-DPC-V2.

#### REFERENCES CITED

- Agnesi, V., Camarda, M., Conoscenti, C., Di Maggio, C., Diliberto, I.S., Madonia, P., and Rotigliano, E., 2005, A multidisciplinary approach to the evaluation of the mechanism that triggered the Cerda landslide (Sicily, Italy): *Geomorphology*, v. 65, p. 101–116, doi: 10.1016/j.geomorph.2004.08.003.
- Allard, P., Behncke, B., D'Amico, S., Neri, M., and Gambino, S., 2006, Mount Etna 1993–2005: Anatomy of an evolving eruptive cycle: *Earth-Science Reviews*, v. 78, p. 85–114, doi: 10.1016/j.earscirev.2006.04.002.
- Billi, A., and Funicello, R., 2008, Concurrent eruptions at Etna, Stromboli, and Vulcano: Casualty or causality?: *Annales de Geophysique*, v. 51 (in press).
- Billi, A., Presti, D., Faccenna, C., Neri, G., and Orecchio, B., 2007, Seismotectonics of the Nubia plate compressive margin in the south Tyrrhenian region, Italy: Clues for subduction inception: *Journal of Geophysical Research*, v. 112, B08302, doi: 10.1029/2006JB004837.
- Boore, D.M., 1983, Stochastic simulation of high-frequency ground motions based on seismological models of the radiated spectra: *Bulletin of the Seismological Society of America*, v. 73, p. 1865–1894.
- Brodsky, E.E., and Prejean, S.G., 2005, New constraints on mechanisms of remotely triggered seismicity at Long Valley Caldera: *Journal of Geophysical Research*, v. 110, p. 1–14.
- Brune, J.N., 1970, Tectonic stress and the spectra of seismic shear waves from earthquakes: *Journal of Geophysical Research*, v. 75, p. 4997–5009, doi: 10.1029/JB075i026p04997.
- Capaccioni, B., Tassi, F., Vaselli, O., Tedesco, D., and Poreda, R., 2007, Submarine gas burst at Panarea Island (southern Italy) on 3 November 2002: A magmatic versus hydrothermal episode: *Journal of Geophysical Research*, v. 112, B05201, doi: 10.1029/2006JB004359.
- Caracausi, A., Di Alberto, V., and Grassa, F., 2002, Effetti del sisma del 6/9/2002 sulle sorgenti termali della Sicilia Nord Occidentale: Palermo, Istituto Nazionale di Geofisica e Vulcanologia (INGV) Open File Report: <http://www.pa.ingv.it/Sisma2002/report.htm> (July 2008).
- Cigolini, C., Laiolo, M., and Coppola, D., 2007, Earthquake-volcano interactions detected from radon degassing at Stromboli (Italy): *Earth and Planetary Science Letters*, v. 257, p. 511–525, doi: 10.1016/j.epsl.2007.03.022.
- D'Agostino, N., and Selvaggi, G., 2004, Crustal motion along the Eurasia-Nubia plate boundary in the Calabrian Arc and Sicily and active extension in the Messina Straits from GPS measurements: *Journal of Geophysical Research*, v. 109, B11402, doi: 10.1029/2004JB002998.
- De Astis, G., Ventura, G., and Vilardo, G., 2003, Geodynamic significance of the Aeolian volcanism (southern Tyrrhenian Sea, Italy) in light of structural, seismological, and geochemical data: *Tectonics*, v. 22, 1040, doi: 10.1029/2003TC001506.
- Esposito, A., Giordano, F., and Anzidei, M., 2006, The 2002–2003 submarine gas eruption at Panarea volcano (Aeolian Islands, Italy): *Volcanology of the seafloor and implications for the hazard scenario: Marine Geology*, v. 227, p. 119–134, doi: 10.1016/j.margeo.2005.11.007.
- Feuillet, N., Cocco, M., Musumeci, C., and Nostro, C., 2006, Stress interaction between seismic and volcanic activity at Mt. Etna: *Geophysical Journal International*, v. 164, p. 697–718, doi: 10.1111/j.1365-246X.2005.02824.x.
- Gvirtzman, Z., and Nur, A., 1999, The formation of Mount Etna as the consequence of slab rollback: *Nature*, v. 401, p. 782–785, doi: 10.1038/44555.
- Hill, D.P., 2008, Dynamic stresses, coulomb failure, and remote triggering: *Bulletin of the Seismological Society of America*, v. 98, p. 66–92, doi: 10.1785/0120070049.
- Lanzafame, G., and Bousquet, J.C., 1997, The Maltese escarpment and its extension from Mt. Etna to Aeolian islands (Sicily): Importance and evolution of a lithospheric discontinuity: *Acta Vulcanologica*, v. 9, p. 121–135.
- Linde, A.T., and Sacks, I.S., 1998, Triggering of volcanic eruptions: *Nature*, v. 395, p. 888–890, doi: 10.1038/27650.
- Lucchi, F., Tranne, C.A., Calanchi, N., and Rossi, P.L., 2006, Late Quaternary deformation history of the volcanic edifice of Panarea, Aeolian Arc, Italy: *Bulletin of Volcanology*, v. 69, p. 239–257, doi: 10.1007/s00445-006-0070-9.
- Manga, M., and Brodsky, E.E., 2006, Seismic triggering of eruptions in the far field: *Volcanoes and geysers: Annual Review of Earth and Planetary Sciences*, v. 34, p. 263–291, doi: 10.1146/annurev.earth.34.031405.125125.
- Neri, G., Caccamo, D., Cocina, O., and Montalto, A., 1996, Geodynamics implications of the earthquake data in the southern Tyrrhenian sea: *Tectonophysics*, v. 258, p. 233–249, doi: 10.1016/0040-1951(95)00202-2.
- Neri, M., Accocella, V., Behncke, B., Maiolino, V., Ursino, A., and Velardita, R., 2005, Contrasting triggering mechanisms of the 2001 and 2002–2003 eruptions of Mount Etna (Italy): *Journal of Volcanology and Geothermal Research*, v. 144, p. 235–255, doi: 10.1016/j.jvolgeores.2004.11.025.
- Ripepe, M., Marchetti, E., Olivieri, G., Harris, A., Dehn, J., Burton, M., Caltabiano, T., and Salerno, G., 2005, Effusive to explosive transition during the 2003 eruption of Stromboli volcano: *Geology*, v. 33, p. 341–344, doi: 10.1130/G21173.1.
- Rovelli, A., Vuan, A., Mele, G., Priolo, E., and Boschi, E., 2004, Rarely observed short-period (5–10 s) suboceanic Rayleigh waves propagating across the Tyrrhenian Sea: *Geophysical Research Letters*, v. 31, L22605, doi: 10.1029/2004GL021194.
- Walter, T.R., and Amelung, F., 2007, Volcanic eruptions following M9 megathrust earthquakes: Implications for the Sumatra-Andaman volcanoes: *Geology*, v. 35, p. 539–542, doi: 10.1130/G23429A.1.
- Wang, R., 1999, A simple orthonormalization method for stable and efficient computation of Green's functions: *Bulletin of the Seismological Society of America*, v. 89, p. 733–741.

Manuscript received 5 August 2008  
 Revised manuscript received 22 October 2008  
 Manuscript accepted 26 October 2008

Printed in USA